

1 **Conflicting edge influence on herbaceous species in open areas vs. underneath oak trees in**
2 **forest fragments in Iran**

3 *Gelareh Valadi¹, Javad Eshaghi Rad^{2*}, Yahia Khodakarami³, Karen Amanda Harper⁴*

4 *1-Department of Forestry, Faculty of Natural Resources, Urmia University, Iran.*

5 *2-Corresponding author. Department of Forestry, Faculty of Natural Resources, Urmia*
6 *University, Iran*

7 *3-Forests and Rangelands Research Department, Agricultural research, Education and*
8 *extension organization, AREEO, Kermanshah, Iran*

9 *4-Biology Department, Saint Mary's University, Halifax, Canada*

10 * Correspondence: Javad Eshaghi Rad

11 Email: j.eshagh@urmia.ac.ir or Javad.eshaghi@yahoo.com

12 Tel: +98 44 32770489, Fax: +98 44 32770489

13

14 **Abstract**

15 Since the type of forest influences vegetation patterns from the edge to interior forest, site-specific
16 edge studies are needed but there have been few studies in open-canopied forests such as oak
17 savannahs. Our objective was to compare patterns of herbaceous plant diversity along the forest
18 edge-to-interior gradient between open areas and underneath oak trees in the Zagros Forest in Iran.
19 We established eighteen transects from the forest edge to the interior in small and large forest
20 fragments to sample herbaceous species in five 0.25 m² quadrats at 1 m intervals from the base of
21 the tree to the open area at different distances from the forest edge. We analyzed the data using

22 randomization tests for edge influence and generalized linear mixed models. Edge influence had a
23 positive effect on herbaceous species richness and diversity underneath oak trees but a negative
24 effect in open areas. At forest edges, species richness and diversity significantly decreased from
25 the tree base toward open areas but exhibited the opposite pattern away from the edge. Edge
26 influence extended up to 50 m from the forest edge to the interior. Our findings highlight the
27 importance of considering forest type and stand heterogeneity when studying edge influence on
28 plant diversity. Our results show that edge studies are needed for specific forest types, particularly
29 in heterogeneous landscapes, to ensure appropriate conservation of species diversity. We
30 recommend establishing a 50-meter buffer zone along edges in the Zagros Forest in Iran to
31 minimize negative edge influence on herbaceous plant diversity.

32

33 **Keywords:** Distance of edge influence, edge effects, fragment size, open canopy forest, species
34 richness, Zagros forests.

35

36 **Introduction**

37 Destruction and degradation of natural ecosystems are the primary causes of the decrease
38 in global biodiversity (Rands et al. 2010). Human disturbances such as logging, forest clearing for
39 agriculture, and landscape fragmentation are related to loss of natural habitat and biological
40 diversity (Barima et al. 2010). Fragmentation, the division of natural habitat into smaller and more
41 isolated fragments (Haddad et al. 2015), alters forest dynamics, microclimate, and biological
42 cycles, leading to an increase in invasive and pioneer species (Barima et al. 2010), and changes in

43 environmental factors, community structure, and species composition close to the edge of
44 fragments (Harper 2005; Pardini et al. 2017).

45 One of the major consequences of forest fragmentation is an increase in forested areas
46 influenced by the edge (Honnay et al. 2002; Fahrig 2003). This concept is termed edge influence
47 and is defined as the difference in biotic and abiotic factors at the forest edge relative to the interior
48 forest (Harper et al. 2005). Edge influence can have important impacts on species diversity, and
49 community and ecosystem functioning (Laurance et al. 2006; Willmer et al. 2022). Along a forest
50 edge-to-interior gradient, species are exposed to changes in microclimatic conditions such as
51 greater light availability, temperature variation, and wind exposure (Harper et al. 2005; Magnago
52 et al. 2015; Erdős et al. 2018), which affect the establishment and development of plants (Coelho
53 et al. 2016; Erdős et al. 2019; Wekesa et al. 2019; da Costa et al. 2020). The edge is often
54 dominated by light-demanding species with high growth and low survival rates (Magnago et al.
55 2015; Bragion et al. 2019). In contrast, the shady and humid conditions in the forest interior favor
56 long-lived shade-tolerant species (Bragion et al. 2019), which grow slowly but are taller and larger,
57 resulting in greater aboveground stand biomass (Da Silva et al. 2019). Edges also influence litter
58 decomposition and nutrients, and subsequently alter species diversity and richness along the forest
59 edge-to-interior gradient (Bennett and Saunders 2010).

60 Edge influence has been a principal topic of interest in studies of landscape processes
61 associated with edge creation and fragmentation during the last few decades (Harper et al. 2005,
62 Franklin et al. 2021). Forest herbaceous species can be influenced by the edge because their
63 composition is affected by altered forest conditions such as increased light availability and reduced
64 soil moisture (Pellissier et al. 2013). Furthermore, conditions at the forest edge have been found

65 to be more heterogeneous compared to the interior (Ewers and Didham 2006). Previous studies
66 have shown that forest edges influence woody plant species richness and diversity in different
67 ecosystems including in South Africa (Ruwanza 2018), Tanzania (Kacholi 2014) and Brazil
68 (Fontoura et al. 2006; Sampaio and Scariot 2011). However, few studies have investigated the
69 herbaceous layer diversity in response to created edges. Studies show that plant species richness
70 and diversity of understory species decreased from the forest edge to interior forest in central-
71 southern China (Li et al. 2018) and in southwestern France (Alignier 2013). However, the opposite
72 trend of higher species richness in the forest interior compared to edge has been found in northern
73 France and Atlantic Forest in Brazil (Berges et al. 2013; Mendes et al. 2016). Finally, no edge
74 influence on species richness was reported in southwestern Amazon forests (Phillips et al. 2006).

75 Furthermore, studies of edge influence on vegetation in open, dry forests are compared to
76 those in more humid ecosystems. For instance, studies have been conducted in humid black spruce
77 boreal forests in Canada (Harper et al. 2016) and the tropical cerrados in Brazil (Dodonov et al.
78 2013). Moreover, no edge research has considered differences in edge influence on herbaceous
79 vegetation in different habitats within a heterogeneous open-canopied forest or the interaction
80 between edge influence (forest edge-to-interior gradient) and the gradient from the tree base to
81 open area away from the tree canopy.

82 The Zagros Forest, an open-canopied temperate forest dominated by *Quercus* spp., is the
83 largest forested land in Iran and has been fragmented by human activities such as fuelwood cutting,
84 agriculture and livestock grazing. The forests have been significantly destroyed and their potential
85 productivity has been lost due to social problems and inadequate management practices (Eshaghi
86 Rad et al. 2018). In a previous study in this forest, we investigated edge influence on herbaceous

87 plant species diversity and soil properties along the forest edge-to-interior gradient (Valadi et al.
88 2022). Here we investigate edge influence further by considering the effect of distance from the
89 tree base into an open area on herbaceous species richness and diversity at different distances along
90 the forest edge-to-interior gradient. Our first objective was to determine how herbaceous plant
91 richness and diversity varied along two gradients: (i) from the tree base to open area and (ii) from
92 the forest edge to the interior, and to ascertain whether these two gradients interact. Our second
93 objective was to assess the differences in herbaceous plant richness and diversity in small vs. large
94 fragments. We tested the following null hypotheses: (i) species richness and diversity is the same
95 at different distances from the base of tree, (ii) changes in species richness and diversity from the
96 forest edge to interior are the same at different distances from the tree base toward open area, and
97 (iii) herbaceous plant richness and diversity patterns along gradients are the same in small and
98 large forest fragments. By understanding the effects of edge influence on herbaceous species
99 richness and diversity, forest managers could develop more effective strategies to conserve and
100 protect these important ecosystems.

101

102 **Material and Methods**

103 **Study area**

104 We conducted our research in the semi-arid Kermanshah province in Iran (34°1'20.37" N,
105 46°23'54.93"E, 1650 m asl). *Quercus brantti*, the main tree species in our study area, forms even-
106 aged stands with a density of 70 individuals per ha and canopy cover < 50% (Jazirei and Ebrahimi
107 Rastaghi 2003). Average annual precipitation and temperature were 489 mm and 21.4 °C,
108 respectively. The lowest and highest monthly average temperatures were 8.2 °C in January and

109 35.2 °C in August 2019. From the past to present, these forests have been settled by residents and
110 nomads resulting in deforestation in some parts and severe damages in others. Due to the lack of
111 adequate conservation planning, this settlement created forest fragments of varying sizes.

112

113 **Data collection**

114 To investigate edge influence on species richness and diversity of herbaceous vegetation
115 in sparse oak forests, we selected three small (5 to 7 ha) and three large (13 to 18 ha) fragments on
116 20-25% north-facing slopes. We chose fragments with similar physiographical conditions to
117 isolate the effect of edge influence and we maintained a distance of approximately 1 km between
118 fragments. We established three transects from the edge to the forest interior in each of the three
119 small and three large forest fragments for a total of 18 transects. The first transect in each fragment
120 was randomly chosen (using random coordinates) and the other two transects were placed 200 m
121 on either side parallel to the first one. Herbaceous vegetation was sampled in May and June 2019
122 at 0 (forest edge), 25, 50, 100, and 150 m distances (toward forest interior) along each transect
123 (Mendes et al. 2016) for a total of 90 sampling points in the six forest fragments (15 per fragment,
124 45 in small and 45 in large fragments).

125 To understand how herbaceous vegetation richness and diversity change from the tree base
126 to the adjacent open area we collected data on canopy cover. We measured the short and long
127 crown diameters of all trees with a DBH greater than 7.5 cm in two quadrats (20 × 2 m)
128 perpendicular to the main edge to interior transect at each sampling point. We collected herbaceous
129 data in five 0.5 × 0.5 m (0.25 m²) quadrats at 1 m intervals from the base of two trees at each
130 sampling point (ten quadrats total). We selected the nearest tree on either side of the main transect

131 and established the quadrats from the tree base towards open area and the main transect (Fig.1).
132 We recorded the number of individuals of all vascular herbaceous species < 0.5 m in height within
133 each quadrat. Individuals were easily differentiated for most species, but we estimated the number
134 of individuals for a few species with high density such as some grasses. Herbaceous species were
135 identified to species level; nomenclature followed Ghahraman (2001).

136

137 **Data analysis**

138 For each sampling point, we calculated the mean herbaceous species abundance for paired
139 quadrats located at the same distance from the tree base for a total of five mean abundances (one
140 for each distance from the tree base) for each species at each sampling point. Before data analysis,
141 Shapiro-Wilk tests were used to test for data normality. Unless otherwise indicated, all data
142 analyses were conducted in R version 3.6.1 (R Core Team 2014).

143 Herbaceous vegetation diversity was quantified for each sampling point using three
144 diversity indices: species richness (N = number of species), Shannon diversity as $H' =$
145 $\sum_{i=1}^s p_i \ln p_i$, where s equals the number of species and p_i is the relative cover of i^{th} species
146 (hereafter referred to as diversity) and evenness as $J' = H' / H'_{max}$ with $H'_{max} = \ln(S)$ (Magurran
147 2004). We analyzed diversity using the package “vegan,” version 2.5-6 (Oksanen et al. 2013). We
148 calculated average canopy cover by using $CD = (C1 \times C2) \times \pi / 4$ where CD =canopy diameter,
149 $C1$ =long diameter, $C2$ = short diameter (Zobeyri 2008) for each tree, which we then averaged for
150 all sampling points for each transect. We detected significant differences in canopy cover between
151 different distance from edge using the Tukey test in SPSS 22 (Rovai et al. 2013).

152 For each of the five distances from the tree base, we calculated the magnitude of edge
153 influence (MEI) and distance of edge influence (DEI) (Harper and Macdonald 2011) for species
154 richness, diversity and evenness. The MEI is a measure of the strength of edge influence, which
155 we determined as $MEI = (X_d - X_i)/(X_d + X_i)$ where X_d = average of each variable at distance d
156 from the edge, and X_i = average of each variable in interior forest (100 m and 150 m). This metric
157 ranges from -1 (negative edge influence) to +1 (positive edge influence). We reported MEI at the
158 distance from the edge where the absolute value of MEI was greatest for each variable. To calculate
159 DEI for each variable, we used the randomization test of edge influence (RTEI) according to the
160 methodology in Harper et al. (2011). RTEI tests the significance of MEI for various distances from
161 the edge compared to interior forest using randomization tests of the data. We reported DEI as
162 either 0 m if MEI was significant only at 0 m or the set of two or more consecutive distances (or
163 separated by one distance) where MEI was significant. Otherwise, DEI was reported as not
164 significant and was excluded from average DEI. We calculated MEI and DEI separately for the
165 five distances from the tree base into the open area.

166 We used generalized linear mixed models (GLMMs) (Magnago et al. 2017) to assess the
167 effects and interactions of distance from forest edge, distance from tree base and fragment size on
168 diversity indices. Distance from edge, distance from tree base and fragment size were fixed effects
169 and fragment was a random effect. A Gaussian distribution was used for the normally distributed
170 response variables. For analyzing GLMMs, we used the package “lme4” version 1.1-21 (Bates et
171 al. 2014). Tukey tests (in SPSS 22) were used to compare diversity indices at different distances
172 from the edge for each distance from the tree base (Rovai et al. 2013). Indicator species analysis
173 was applied to determine indicator species for different distances from the tree base in small and

174 large fragments (McCune and Mefford 2006). This method is based on relative fidelity and relative
175 abundance of species and aims to identify species (Legendre and Legendre 2012).

176

177 **Results**

178 Trees had significantly larger canopies in the forest interior compared to the edge in both
179 small and large fragments; canopy area per tree was particularly low within 50 m of the edges of
180 large fragments (Table 1).

181 The results of the GLMM showed that distance from forest edge and distance from tree
182 base significantly affected herbaceous plant species richness, diversity and evenness (Table 2).
183 Furthermore, the interaction between distance from edge and distance from tree base was
184 significant. Fragment size was a significant factor in explaining species diversity and evenness,
185 but not richness. The interactions of fragment size with distance from edge and with distance from
186 tree base were significant except for the interaction between fragment size and distance from edge
187 for species diversity, and the interaction between fragment size and distance from tree base for
188 species diversity and evenness.

189 At the edges of small and large forest fragments (0, 25 m), herbaceous plant species
190 richness and diversity significantly decreased from the tree base (0, 1, 2 m) toward open area (3,
191 4 m) (Fig. 2). We found the opposite pattern in interior forest, with significantly higher species
192 richness and diversity 3 and 4 m from the tree base. Evenness was significantly greater in the open
193 area than at the base of trees at distances of 150 m from the edge in small fragments, and 100 m
194 and 150 m from the edge in large fragments. The interaction between distance from tree base and
195 distance from forest edge can also be viewed from a different perspective. Measures of species

196 diversity at the tree base decreased from the forest edge to the interior but increased along the
197 edge-to-interior gradient in open areas. Overall, diversity was lowest in open areas near the edge
198 and next to tree bases in the forest interior, and greatest at tree bases at the edge and in open areas
199 of interior forest.

200 The MEI was positive (greater values at the edge) for herbaceous plant species richness,
201 diversity and evenness for areas within 3 m of the tree base in both small and large fragments, but
202 negative for distances greater than 3 m from the tree base in the open areas (Table 3). The DEI for
203 species richness and diversity extended up to 50 m from the forest edge to the interior for nearly
204 all distances from the tree base in both small and large forest fragments.

205 Herbaceous plant indicator species were discernible only for the tree base (0, 1 m) at the
206 forest edge (0 m) and in open areas (4 m from the tree base) at 100 and 150 m from the edge in
207 small forest fragments (Table 5). For large fragments, indicator species were identified for
208 comparable distances from the tree base and the forest edge, with the addition of the tree base (0,
209 1 m) at a distance of 25 m from the edge.

210

211 **Discussion**

212 Overall, we found opposing patterns of edge influence on herbaceous understory
213 vegetation in oak savannah forest fragments in the Zargos Forest of Iran. Edge influence was
214 positive for herbaceous plant species diversity at tree bases but negative in the open areas between
215 trees (Table 3). Stated another way, diversity was greater under trees than in open areas up to 50
216 m from the edge, but the opposite pattern occurred in the forest interior (100 and 150 m from the
217 forest edge) with greater diversity in open areas. Microenvironmental variation along the forest

218 edge-to-interior gradient might explain these opposing patterns; a different microclimate at the
219 edge may favor a different plant community from that found in the interior (Noss and Cooperrider
220 1994). Documented changes in microclimate typical of forest edges include higher light, air and
221 soil temperatures, wind speed, and vapor pressure, and lower relative humidity and soil moisture
222 (Young and Mitchell 1994). Increased evaporation and reduced soil moisture adjacent to the forest
223 edge are crucial drivers behind differences in forest vegetation between forest edge and interior
224 (Herbst et al. 2007; Riutta et al. 2016). Soil carbon and moisture levels are higher in shaded areas
225 than in open areas at the forest edge (Joshi et al. 2001). Combined with additional light penetration
226 and more organic matter, these wetter conditions under the canopy at the forest edge likely favor
227 more species, resulting in higher richness and diversity compared to the drier, nutrient-poor
228 conditions in open areas. Greater herbaceous species richness under tree canopies near the forest
229 edge is associated with more organic matter and soil moisture, wind protection, decreased daily
230 oscillations of temperature, and lower evapotranspiration rates, air, and soil temperatures (Ishii
231 2013; Valladares 2016; Ren 2022).

232 Edge influence did not affect herbaceous species richness and diversity after 50 m. In
233 contrast to forest edges, interior forest had greater canopy cover (Table 1), resulting in less light
234 availability. Although soil moisture is generally important, light is probably the most limiting
235 factor for understory species in temperate forests (Dormann et al. 2020). This lack of light is more
236 important for the establishment of herbaceous species, as shade reduces herbaceous species
237 richness (Gillet et al. 1999; Fikadu and Zewdu 2021). Light is a key resource for the growth and
238 survival of herbaceous species (Tinya 2009; Plue et al. 2013; Garg 2022) and is likely the reason
239 we observed more herbaceous species in open areas compared to tree bases in interior forests,

240 which had less available light because of greater canopy cover. Many studies found that light
241 availability has a major impact on herbaceous species composition (e.g., De Frenne et al. 2015;
242 Medvecká et al. 2018). Most herbaceous species in sparse oak forests, such as *Tortilis* sp.,
243 *Hordeum* sp., and *Heterantheium* sp. in open areas within the forest interior, and *Astragalus* sp.
244 and *Trifolium* at forest edges, which mainly belong to Poaceae and Fabaceae, are adapted to high
245 light conditions and are not usually found in low light conditions beneath the canopy. Greater light
246 availability in open-canopied forest tends to promote the establishment of generalist and light-
247 demanding species (Alignier et al. 2014).

248 Distance from the forest edge and from the tree base were crucial factors in the open canopy
249 oak forests, as we found opposite patterns of edge influence on herbaceous species diversity for
250 the tree base vs. open areas (Fig.2). We believe that these results are related to increasing canopy
251 cover from the forest edge to interior, which mediates harsh abiotic environmental conditions such
252 as light availability, wind speed, air temperature, and humidity and reduced soil evaporation (Sagar
253 et al. 2012; Ishii et al. 2013; Valladares et al. 2016). Whereas light availability is positively
254 correlated with understory plant species richness in temperate forests (Dormann 2020), this
255 relationship is not consistent across all forests. Studies have found varying relationships between
256 light availability and plant species richness (Adler 2011; Bartels and Chen 2013; Fuxai et al. 2014;
257 Tinya 2016). These relationships often depend on factors such as dominant tree species, stand
258 density, soil properties, successional stage, and management (Hardtle 2003; Fuxai et al. 2014).
259 Carefully controlled grazing can increase plant diversity (Kirk et al. 2019); a study of Zagros
260 forests showed that herbaceous and woody communities responded differently to various levels of

261 grazing intensity (Ahmadi et al. 2022). In our study fragments were surrounded by agricultural
262 land in which cattle grazing was prohibited by landowners.

263 Our result of a DEI of 50 m for herbaceous species richness and diversity agrees with other
264 studies that indicate that DEI usually extends up to 50 m in temperate forests (Honday et al. 2002)
265 and 40 m in boreal forests (Harper and Macdonald 2001). Based on a synthesis by Franklin et al.
266 (2021), average DEI for forest fragments surrounded by anthropogenic disturbances extends up to
267 42 m into the interior. Guirado et al. (2006) observed greater DEI (100 m) in oak and pine
268 Mediterranean forests in Spain, indicating that DEI depends on various conditions in different
269 ecosystems. Forest practices can strongly modify understory environmental conditions such as
270 light, temperature, and soil moisture as well as species diversity (Ash and Barkham 1976; Grayson
271 et al. 2012). In the Zagros Forest in Iran, considering that DEI extended up to 50 m for both under
272 trees and in open areas, we recommend a 50 m buffer to conserve the interior herbaceous
273 communities of these oak fragments. Further research on the impact of edge influence and buffer
274 zones is urgently required in these fragmented forests to develop comprehensive management
275 plans for each forest.

276 In conclusion, our study showed that forest edges influence herbaceous species richness
277 and diversity and have different impacts on species at the tree base compared to in open areas in
278 open-canopied oak forests of Iran. Efforts to conserve and restore forests and herbaceous plants
279 should be integrated with sustainable forest management practices to maintain and enhance the
280 ecosystem services of these forests, ensuring their benefits for present and future generations.
281 Regional and national assessments are needed to determine where and what kind of conservation
282 and restoration should occur to protect the remaining natural herbaceous plants. Our study has

283 major implications for edge research beyond open oak forests in that we showed how edge
284 influence on plant diversity can differ dramatically at a fine scale within the same ecosystem, even
285 having opposite effects within a few meters from the tree base.

286

287 **References**

288 Adler P B, Seabloom E W, Borer E T et al. (2011) Productivity is a poor predictor of plant
289 species richness. *Science* 333(6050):1750-1753

290 Ahmadi A, Rezaei R, Abdi N, Toranjzar H (2022) Changes in species diversity and soil seed
291 bank under the enclosure and different intensity of livestock grazing in deteriorated forests of
292 middle Zagros. *Ecol Iran For* 10(20): 52-63

293 Alignier A, Deconchat M (2013) Patterns of forest vegetation responses to edge effect as
294 revealed by a continuous approach. *Ann For Sci* 70(6): 601-609

295 Alignier A, Alard D, Chevalier R, Corcket E (2014) Can contrast between forest and adjacent
296 open habitat explain the edge effects on plant diversity? *Acta Bot Gall* 161(3): 253-259

297 Ash J, Barkham J (1976) Changes and variability in the field layer of a coppiced woodland in
298 Norfolk, England. *J. Ecol* 697-712

299 Barima Y S S, Barbier N, Ouattara B, Bogaert J (2010) Relation entre la composition floristique
300 et des indicateurs de la fragmentation du paysage dans une region de transition forêt savane

301 ivoirienne [Relationship between floristic composition and indicators of landscape fragmentation
302 in a transition forest-savanna region in Ivory Coast]. *Biotechnol Agron Soc Envir* 14(4): 617

303 Bartels S F, Chen HYH (2010) Is understory plant species diversity driven by resource quantity
304 or resource heterogeneity? *Ecol* 91(7):1931-1938

305 Bates D, Mächler M, Bolker B et al (2014) Fitting linear mixed-effects models using lme4.

306 ArXiv org :1406.5823

307 Bennett A F, Saunders D A (2010) Habitat fragmentation and landscape change. *Conserv Biol*
308 93: 1544-1550

309 Bergès L, Chevalier R, Avon C (2013) Influence of forest road, road-surfacing material and
310 stand age on floristic diversity and composition in a nutrient-poor environment. *Appl Veg Sci*
311 16: 470– 479

312 Bragion E D F A, Coelho G A O, de Siqueira F F et al (2019) Sharp differentiation on the
313 performance of plant functional groups across natural edges. *Plant Ecol* 12(1): 186-198

314 Coelho R L F, de Oliveira D S, de Almeida M I S (2016) Does social media matter for post
315 typology? Impact of post content on Facebook and Instagram metrics. *Online Inf Rev* 40(4):
316 458-471

317 Da Costa L B, de Carvalho O L F, de Albuquerque A O et al (2022) Deep semantic segmentation
318 for detecting eucalyptus planted forests in the Brazilian territory using sentinel-2 imagery.
319 *Geocarto Int* 37(22): 6538-6550

320 Da Silva T S, Haigh S K, Elshafie M Z E B et al (2019) Preparations for field testing for the
321 performance validation of piled wind turbine foundations in expansive clays. *ISSMGE*

322 De Frenne P, Rodríguez-Sánchez F, De Schrijver A N (2015) Light accelerates plant responses
323 to warming. *Nat Plants* 1: 1-3. doi:10.1038/nplants.2015.110

324 Dodonov P, Harper K A, Silva-Matos D M (2013) The role of edge contrast and forest structure
325 in edge influence: vegetation and microclimate at edges in the Brazilian cerrado. *Plant Ecol*
326 214(11): 1345-1359

327 Dormann C F, Bagnara M, Boch S et al (2020) Plant species richness increases with light
328 availability, but not variability, in temperate forests understorey. *BMC Ecol* 20: 1-9

329 Erdős L, Ambarlı D, Anenkhonov O A et al (2018) The edge of two worlds: A new review and
330 synthesis on Eurasian forest-steppes. *Appl Veg Sci* 21 (3): 345-362

331 Erdős L, Krstonošić D, Kiss PJ et al (2019) Plant composition and diversity at edges in a semi
332 natural forest–grassland mosaic. *Plant Ecol* 220(3):279-292

333 Eshaghi Rad J, Valadi G, Salehzade O, Maroufi H (2018) Effects of anthropogenic disturbance
334 on
335 plant composition plant diversity and soil properties in oak forests, Iran. *J For Sci* 64(8): 358-370

336 Ewers R M, Didham R K (2006) Confounding factors in the detection of species responses to
337 habitat fragmentation. *Biol Rev* 81:117-707

338 Fahrig L (2003) Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol Syst*
339 34(1): 487-515

340 Fikadu T, Zewdu T (2021) Influences of tree species and canopy cover on aboveground Biomass
341 yield and ground cover of herbaceous plants in eastern Oromia, Ethiopia. *Am J Agric For* 9(4):
342 233-240

343 Fontoura S B, Ganade G, Larocca J (2006) Changes in plant community diversity and
344 composition across an edge between *Araucaria* forest and pasture in South Brazil. *Am J Bot* 29:
345 79-91

346 Franklin C M A, Harper K A, Clarke M J (2021) Trends in studies of edge influence on
347 vegetation at human-created and natural forest edges across time and space. *Can J For Res* 51:
348 274-282

349 Fuxai X, Fousseni F, Chungang P et al (2014) Effect of overstory on the seasonal variability of
350 understory herbs in primary broad-leaved Korean pine forest of Changbai Mountain.
351 *AJB* 13(11). <https://doi.org/10.5897/ajb12.1013>

352 Garg S, Joshi R K, Garkoti S C (2022) Effect of tree canopy on herbaceous vegetation and soil
353 characteristics in semi-arid forests of the Aravalli hills. *Arid Land Res Manag* 36(2): 224-242

354 Ghahreman A (1979-2003) *Color flora of Iran*. Research Institute of Forests and Rangelands,
355 Tehran

356 Gillet F B, Murisier A, Buttler J, Gallandat J (1999) Influence of tree cover on the diversity of
357 herbaceous communities in subalpine wooded pastures. *Appl Veg Sci* 2: 47-54

358 Grayson S F, Buckley D S, Henning J G et al (2012) Understory light regimes following
359 silvicultural treatments in central hardwood forests in Kentucky, USA. *For Ecol Manag* 279: 66-
360 76

361 Guirado M, Pino J, Rodà F (2006) Understorey plant species richness and composition in
362 metropolitan forest archipelagos: effects of forest size, adjacent land use and distance to the
363 edge. *Glob Ecol Biogeogr* 15(1): 50-62

364 Haddad N M, Brudvig L A, Clobert J et al (2015) Habitat fragmentation and its lasting impact on
365 Earth's ecosystems. *Sci Adv* 1(2) e1500052

366 Härdtle W, von Oheimb G, Westphal C (2003) The effects of light and soil conditions on the
367 species richness of the ground vegetation of deciduous forests in northern Germany (Schleswig-
368 Holstein). For Ecol Manag 182:327–38

369 Harper K A, Macdonald S E (2001) Structure and composition of riparian boreal forest: new
370 methods for analyzing edge influence. Ecol 82(3):649-659

371 Harper K A, Macdonald S E, Burton P J (2005) Edge influence on forest structure and
372 composition in fragmented landscapes. Conserv Biol 19(3): 768-782

373 Harper K A, Macdonald S E (2011) Quantifying distance of edge influence: a comparison of
374 methods and a new randomization method. Ecosphere 2(8): 1-17

375 Harper K A, Drapeau P, Lesieur D, Bergeron Y (2016) Negligible structural development and
376 edge influence on the understory at 16-17-yr-old clear-cut edges in black spruce forest.
377 Appl Veg Sci 19:462-473

378 Herbst M, Roberts J M, Rosier P T W, Taylor M E, Gowing D J (2007) Edge effects and forest
379 water use: a field study in a mixed deciduous woodland. For Ecol Manag 250:176–86

380 Honnay O, Verheyen K, Hermy M (2002) Permeability of ancient forest edges for weedy plant
381 species invasion. For Ecol Manag 161 (1-3): 109-122

382 Ishii H, Azuma W, Nabeshima E (2013) The need for a canopy perspective to understand the
383 importance of phenotypic plasticity for promoting species coexistence and light-use
384 complementarity in forest ecosystems. Ecol Res 28: 191–198

385 Jazirei M H, Ebrahimi Rastaghi M (2003) Zagros silviculture. Tehran University Press. 560 pp
386 (In Persian).

387 Joshi B, Singh S P, Rawat Y S, Goel D (2001). Facilitative effect of *Coriaria nepalensis* on
388 species diversity and growth of herbs on severely eroded hill slopes. *Curr Sci* 80: 678-682

389 Kacholi D S (2014) Edge-interior disparities in tree species and structural composition of the
390 Kilengwe forest in Morogoro region, Tanzania. *International Scholarly Research Notices in*
391 *Forests of Different Type and Ages. Pol J Ecol* 66(3):239-249

392 Kirk D A, Hébert K, Goldsmith F B (2019) Grazing effects on woody and herbaceous plant
393 biodiversity on a limestone mountain in northern Tunisia. *PeerJ*, 7, e7296

394 Laurance W F, Nascimento H E, Laurance S G (2006) Rain forest fragmentation and the
395 proliferation of successional trees. *Ecol* 87(2): 469-482

396 Legendre P, Legendre L (2012) *Numerical Ecology*. Elsevier

397 Li J, Zhao Ch, Peng Y et al (2018) Edge feffect on tree growth and species diversity in forests of
398 different type and ages. *Pol J Ecol* 66(3):239-249

399 Liepa L, Rendenieks Z, Jansons Ā et al (2020) The persisting influence of edge on vegetation in
400 hemiboreal *Alnus glutinosa* (L.) Gaertn. swamp forest set-asides adjacent to recently disturbed
401 stands. *Forests* 11(10) 1084

402 Magnago L F S, Rocha M F, Meyer L et al (2015) Microclimatic conditions at forest edges have
403 significant impacts on vegetation structure in large Atlantic forest fragments. *Biodivers. Conserv*
404 24: 2305-2318

405 Magnago L F S, Magrach A, Barlow J (2017) Do fragment size and edge effects predict carbon
406 stocks in trees and lianas in tropical forests?. *Funct Ecol* 31(2): 542-552

407 Magurran A E (2004) Measuring biological diversity. *Afr J Aquat Sci*, 29(2): 285-286

408 McCune B, Mefford M (2006) PC-ORD version 5.10: Multivariate Analysis of Ecological Data.
409 MjM Software Gleneden Beach OR USA

410 Medvecká J, Jarolímek I, Hegedušová K (2018) Forest habitat invasions – Who with whom,
411 where and why. *Ecol Manag* 409: 468–478

412 Mendes P G A, Silva M A M, Guerra T N F (2016) Dynamics and edge effect of an Atlantic
413 Forest fragment in Brazil. *Floresta Ambiente* 23(3): 340-349

414 Noss R F, Cooperrider A Y (1994) Saving nature's legacy: protecting and restoring biodiversity.
415 Island Press Washington D C

416 Oksanen J, Blanchet F G, Kindt R et al (2013) Community ecology package. R package version
417 2(0): 321-326

418 Pardini R, Nichols E, Püttker T (2017) Biodiversity response to habitat loss and fragmentation.
419 *Encyclopedia of the Anthropocene* 3: 229-239.

420 Pellissier L, Pinto-Figueroa E, Niculita-Hirzel H et al (2013) Plant species distributions along
421 environmental gradients: do belowground interactions with fungi matter? *Front Plant Sci* 4, 500

422 Phillips O L, Rose S, Mendoza A M et al (2006) Resilience of southwestern Amazon forests to
423 anthropogenic edge effects. *Conserv Biol* 20(6): 1698-1710

424 Plue J, Van Gils B, De Schrijver A (2013) Forest herb layer response to long-term light deficit
425 along a forest developmental series. *Acta Oecol* 53: 63–72

426 Rands M R, Adams W M, Bennun L (2010) Biodiversity conservation: challenges beyond 2010.
427 *Science* 329(5997): 1298-1303

428 R Core Team (2014) R: A Language and Environment for Statistical Computing. Vienna,
429 Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>

430 Ren Y, Guo M, Yin F et al (2022) Tree cover improved the species diversity of understory
431 spontaneous herbs in a small city. *Forests* 13(8): 1310

432 Riutta T, Clack H, Crockatt M, Slade E M (2016) Landscape-scale implications of the edge
433 effect on soil fauna activity in a temperate forest. *Ecosystems* 19: 534–544

434 Rovai A P, Baker J D, Ponton M K (2013) Social science research design and statistics: A
435 practitioner's guide to research methods and IBM SPSS. Water tree Press LLC.

436 Ruwanza S, Gaertner M, Esler K J (2018) Medium-term vegetation recovery after removal of
437 invasive *Eucalyptus camaldulensis* stands along a South African river. *S Afr J Bot S AFR J*
438 *BOT* 119:63-68

439 Sagar R, Pandey A, Singh J S (2012) Composition, species diversity, and biomass of the
440 herbaceous community in dry tropical forest of northern India in relation to soil moisture and
441 light intensity. *The Environmentalist* 32:485-93

442 Sampaio A B, Scariot A (2011) Edge effect on tree diversity, composition and structure in a
443 deciduous dry forest in central Brazil. *Rev. Arvore* 35(5):1121-1134

444 Saunders D A, Hobbs R J, Margules C R (1991) Biological consequences of ecosystem
445 fragmentation: a review. *Conserv. Biol* 5: 18–32

446 Tinya F, Mihók B, Márialigeti S, Mag Z S, Ódor P (2009) A comparison of three
447 indirect methods for estimating understory light at different spatial scales in temperate mixed
448 forests. *Community Ecol* 10: 81–90

449 Tinya, F, Ódor P (2016) Congruence of the spatial pattern of light and understory vegetation in
450 an old-growth, temperate mixed forest. *For Ecol Manag* 381:84-92

451 Valadi G, Eshaghi Rad J, Khodakarami Y et al (2022) Edge influence on herbaceous plant
452 species, diversity and soil properties in sparse oak forest fragments in Iran. *J. Plant Ecol* 15(2)
453 :413-424

454 Valladares F, Laanisto L, Niinemets U, Zavala M A (2016) Shedding light on shade: ecological
455 perspectives of understorey plant life. *Plant Ecol Divers* 9 (3): 237-251

456 Wekesa C, Kirui B K, Marangan E K, Muturi G M (2019) Variation in forest structure, tree
457 species diversity and above-ground biomass in edges to interior cores of fragmented forest
458 patches of Taita Hills, Kenya. *For Ecol Manag* 440: 48-60

459 Willmer J N G, Puettker T, Prevedello J A (2022) Global impacts of edge effects on species
460 richness. *Biol Conserv* 272, 109654

461 Young A, Mitchell N (1994) Microclimate and vegetation edge effects in a fragmented
462 podocarp-broadleaf forest in New Zealand. *Biol Conserv* 67: 63–72

463 Zobeyri M 2008. *Biometry*, Tehran university. 450pp

464

465 **Statements & Declarations**

466 **Funding**

467 This research was financially supported by the vice chancellor for research and technology of
468 Urmia University and received no external funding.

469

470 **Competing Interests**

471 The authors have no relevant financial or nonfinancial interests to disclose.

472

473 **Author Contributions**

474 All authors contributed to the study conception and design. Data collection and analysis were
475 performed by Gelareh Valadi, Javad Eshaghi Rad, Yahia Khodakarami and Karen Amanda
476 Harper. The first draft of the manuscript was written by Gelareh Valadi and all authors
477 commented on previous versions of the manuscript. All authors read and approved the final
478 manuscript.

479

480

481

482 Table 1. Average canopy area per tree (m²) at different distances from the forest edge in small and
483 large forest fragments. Values at different distances within small or large fragments that share the
484 same letter were not significantly different according to Tukey tests.

485

Distance (m)	Small fragments	Large fragments
0	9.52 ± 3.17 ^b	7.11 ± 0.74 ^b
25	12.00±1.72 ^b	7.6±0.74 ^b
50	11.60±1.04 ^{ab}	9.40±0.91 ^{ab}
100	18.07±1.50 ^a	18.35±3.97 ^a
150	14.40±1.89 ^a	18.67±3.65 ^a

489

490

491

492

493

494

495 **Table 2.** Results of the generalized linear mixed models (GLMMs) showing the effects of
 496 distance from edge, distance from tree base and forest fragment size on species diversity indices.

		Sum of squares	df	Mean square	F	Sig
Species richness	Intercept	28960.22	1	28960.22	5156.12	0.00
	Size	3.92	1	3.92	0.69	0.40
	Edge	921.71	4	230.43	41.02	0.00
	Tree base	625.33	4	158.83	28.27	0.00
	Size * Edge	126.45	4	31.62	5.63	0.01
	Size * Tree base	89.45	4	33.36	3.98	0.00
	Edge * Tree base	9689.69	16	586.79	104.48	0.00
	Size * Edge * Tree base	387.56	16	24.22	4.31	0.00
	Error	2246.67	400	5.62		
Shannon diversity	Intercept	1034.12	1	1034.12	7535.54	0.00
	Size	1.48	1	1.48	10.38	0.01
	Edge	11.64	4	2.91	20.45	0.00
	Tree base	4.98	4	1.24	8.74	0.00
	Size * Edge	1.07	4	0.27	1.87	0.11
	Size * Tree base	0.15	4	0.04	0.27	0.89
	Edge * Tree base	167.73	116	11.04	77.60	0.00
	Size * Edge * Tree base	10.56	16	0.66	4.64	0.00
	Error	56.94	400	0.14		
Evenness	Intercept	223.33	1	224.33	7882.87	0.00
	Size	0.65	1	0.65	18.61	0.00
	Edge	1.97	4	0.49	14.12	0.00
	Tree base	1.27	4	0.34	9.07	0.00
	Size * Edge	0.39	4	0.09	2.78	0.02
	Size * Tree base	0.26	4	0.06	1.86	0.11
	Edge * Tree base	8.39	116	0.52	15.01	0.00
	Size * Edge * Tree base	1.23	16	0.08	2.20	0.00
	Error	19.97	400	0.03		

497 edge= distance from edge, tree base= distance from base of tree

498

499

500

501

502

503

504

505

506

507

508 **Table 3.** Magnitude (MEI) and distance of edge influence (DEI) of species diversity indices for
 509 different distances from the base of tree in small and large forest fragments

510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522

	Distance from tree base (m)	Small fragments		Large fragments	
		MEI	DEI (m)	MEI	DEI (m)
Species richness	0	0.6306	0-50	0.8980	0-50
	1	0.5834	0-50	0.8491	0-50
	2	0.2694	0-50	0.4365	0-50
	3	-0.3346	0-50	-0.4237	0-50
	4	-0.6006	0-50	-0.7852	0-50
Shannon diversity	0	0.4743	0-50	0.8708	0-50
	1	0.4306	0-50	0.7508	0-50
	2	0.1616	0-50	-0.0421	NA
	3	-0.2183	0-50	-0.2897	0-50
	4	-0.4311	0-50	-0.5989	0-50
Evenness	0	0.1893	0-50	0.1893	0-50
	1	0.1375	0-25	0.3731	0-50
	2	0.0239	NA	0.0938	0-50
	3	-0.476	NA	-0.1117	NA
	4	-0.1570	0-25	-0.2707	0-50

523 Table 4. List of herbaceous indicator species at different distances from the tree base and forest
 524 edge in small and large forest fragments
 525
 526
 527

Large fragments			Small fragments		
Distance from edge (m)	Distance from tree base (m)	Species	Distance from edge (m)	Distance from tree base (m)	Species
0	0	<i>Gladiolus atroviolaceus</i> Boiss	0	0	<i>Alyssum marginatum</i> Steud.ex Boiss
0	0	<i>Coronilla varia</i> L.	0	0	<i>Lallemantia iberica</i> (M.Beib.)Fisch. & C.A.Mey
0	0	<i>Velezia rigida</i> L.	0	0	<i>Euphorbia bupleuroides</i> Diels
0	0	<i>Minuartia hamata</i> (Hausskn.)Mattf	0	0	<i>Aegilops triuncialis</i> L.
0	0	<i>Campanula cecilli</i> Rech.f.&Schiman	0	0	<i>Hordeum glaucum</i> Steud.
0	0	<i>Salvia multicaulis</i> Vahl	0	0	<i>Euphorbia macroclada</i> Boiss
0	0	<i>Arenaria serpyllifolia</i> L.	0	0	<i>Cephalaria syriaca</i> (L.) Schrad.exRoem.&Schult.
0	1	<i>Achillea aleppica</i> DC.	0	1	<i>Ornithogalum brachystachys</i> K.Koch
0	1	<i>Eryngium thyrsoideum</i> Boiss	0	1	<i>Trifolium dasyurum</i> C.Presl
0	1	<i>Aegilops triuncialis</i> L.	0	1	<i>Filago arvensis</i> L.
0	1	<i>Alyssum marginatum</i> Steud.ex Boiss	0	1	<i>Teucrium scordium</i> L.
0	1	<i>Tragopogon longrostris</i> Bisch.	100	4	<i>Phlomis lanceolata</i> Boiss. & Hohen
0	1	<i>Lophochloa phleoides</i> (Vill.)Rchb.	100	4	<i>Tortilis leptophylla</i> L.
0	1	<i>Hordeum bulbosum</i> L.	100	4	<i>Heteranthelium piliferum</i> (Banks & Soland)
0	1	<i>Bromus danthoniae</i> Trin.	100	4	<i>Quercus brantii</i> Lindl.
25	0	<i>phlomis persica</i> Benth	100	4	<i>Lamium amplexicaule</i> L.
25	1	<i>Euphorbia inderiensis</i> Less.ex Kar.&Kir.	150	4	<i>Rosularia elymatica</i> (Boiss.& Hausskn.ex Boiss.
25	1	<i>Astragalus cyclophyllon</i> Beck	150	4	<i>Erodium cicutarium</i> (L.)Lher
25	1	<i>Trifolium scabrum</i> L.	150	4	<i>Hordeum bulbosum</i> L.
50	0	<i>Vulpia myuros</i> (L.)C.C.Gmel.	150	4	<i>Echinaria capitata</i> (L.)Desf

528

529

530


531 Continued Table 4. List of herbaceous indicator species at different distances from the tree base
 532 and forest edge in small and large forest fragments
 533

Large fragments		
Distance from edge	Distance from tree base	Species
100	4	<i>Poa bulbosa</i> L.
100	4	<i>Euphorbia macroclada</i> Boiss
100	4	<i>Muscari neglectum</i> Guss.ex Ten.
100	4	<i>Euphorbia cheiradenia</i> Boiss
100	4	<i>Marrubium astracanicum</i> Jacq.
100	4	<i>Senecio vernalis</i> Waldst. & Kit.
150	4	<i>Fritillaria imperialis</i> L.
150	4	<i>Quercus brantii</i> Lindl.
150	4	<i>Atractylis cancellata</i> L.
150	4	<i>Lamium amplexicaule</i> L.
150	4	<i>Daphne mucronata</i> Royle
150	4	<i>Tortilis leptophylla</i> L.
150	4	<i>Hordeum glaucum</i> Steud.
150	4	<i>Heterantherium piliferum</i> (Banks & Soland)
150	4	<i>Ziziphora capitata</i> L.

534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551

552

553



554

555

556 **Fig.1** Sampling design for the data collection. DT0 to DT4 refer to quadrat locations at 0 to 4 m

557 from the tree base nearest to each sampling point along the main transect

558







559

560

561

562

563



565 **Fig.2** Species richness, Shannon diversity and evenness at different distances from the forest edge
566 and different distances from the tree base in small (a, b, c) and large (d, e, f) forest fragments. For
567 a given distance from the tree base, values at different distances from the forest edge that share the
568 same letter were not significantly different according to Tukey tests
569